

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/07, 31/20	A1	(11) International Publication Number: WO 99/09969 (43) International Publication Date: 4 March 1999 (04.03.99)
(21) International Application Number: PCT/EP98/05236 (22) International Filing Date: 18 August 1998 (18.08.98) (30) Priority Data: 97114651.9 23 August 1997 (23.08.97) EP (71) Applicant (for all designated States except US): F. HOFFMANN-LA ROCHE AG [CH/CH]; Grenzacherstrasse 124, CH-4070 Basle (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): BOLLAG, Werner [CH/CH]; Mythenstrasse 10, CH-4054 Basle (CH). OTT, Fritz [CH/CH]; Fachstrasse 20, CH-8942 Oberrieden (CH). (74) Agent: KJELLSAA-BERGER, Hanny; Grenzacherstrasse 124, CH-4070 Basle (CH).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: TREATMENT OF CELL-MEDIATED IMMUNE DISEASES (57) Abstract 9-cis retinoic acid and pharmaceutically acceptable salts and pharmaceutically acceptable hydrolyzable esters thereof, 9-cis retinal and pharmaceutically acceptable acetals thereof, and 9-cis retinol and pharmaceutically acceptable hydrolyzable esters thereof as well as metabolites of 9-cis retinoic acid have been found to be efficacious in treating T-helper cell type 1 mediated immune diseases in well tolerated doses. Preferably, the active ingredient is formulated as a medicament for oral or topical administration.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Treatment of Cell-Mediated Immune Diseases

The present invention relates to the use of 9-cis
5 retinoic acid and derivatives or precursors thereof for the
manufacture of a medicament for the treatment of T-helper
cell type 1 mediated immune diseases as well as to the use
of said active substances for the treatment of such dis-
eases.

10 Despite intensive clinical research with retinoids in
the last 27 years, retinoids have not been reported to be
clinically useful in the therapy of immunologically
mediated diseases. Neither diseases caused by T-helper
type-1 cell (Th1) dependent cellular immunity, nor diseases
15 caused by T-helper type-2 cell (Th2) dependent humoral
immunity, have been reported to respond to retinoids. As to
the classification into Th1 dependent diseases - such as
autoimmune and other cell-mediated immune diseases, e.g.
rheumatoid arthritis, multiple sclerosis, uveoretinitis,
20 thyreoiditis, Crohn's disease, insulin dependent diabetes
mellitus, eczema and systemic lupus erythematosus, as well
as rejection of allogeneic organ transplants - and Th2
dependent diseases - i.e. diseases with dominant humoral or
antibody-mediated diseases such as allergic disorders, e.g.
25 atopic dermatitis, allergic rhinitis, hay fever and
allergic bronchial asthma - reference is made to Romagnani,
ed, Th 1 and Th 2 Cells in Health and Disease. Chem.
Immunol., Karger, Basel, 63, pp. 158-170 and 187-203
(1996).

For the first time, quite unexpectedly, it has now been found that a retinoid - namely 9-cis retinoic acid as well as its salts, its esters and its metabolic precursors or prodrugs as well as metabolites of 9-cis retinoic acid, 5 such as 4-oxo-9-cis retinoic acid - is clinically efficacious in the therapy of Th1 dependent diseases.

In the scope of the present invention the term „metabolic precursors and prodrugs as well as metabolites of 9-cis retinoic acid“ encompasses compounds that are converted 10 metabolically into 9-cis retinoic acid, and it includes, in particular, 9-cis retinal and 9-cis retinol as well as pharmaceutically acceptable acetals of 9-cis retinal and pharmaceutically acceptable hydrolyzable esters of 9-cis retinol as well as metabolites of 9-cis retinoic acid such 15 as 4-oxo-9-cis retinoic acid or their glucuronides.

In accordance with this invention, it has thus been found that that administration of 9-cis retinoic acid, its pharmaceutically acceptable salts, its pharmaceutically acceptable hydrolyzable esters, 9-cis retinal, its pharmaceutically acceptable acetals, 9-cis retinol and its pharmaceutically acceptable hydrolyzable esters, as well as 20 metabolites of 9-cis retinoic acid, are efficacious in treating patients with T-helper cell type 1 (Th1) mediated diseases.

25 The invention therefore relates to the use of 9-cis retinoic acid, a pharmaceutically acceptable salt or a pharmaceutically acceptable hydrolyzable ester thereof, 9-cis retinal or a pharmaceutically acceptable acetal thereof or 9-cis retinol or a pharmaceutically acceptable hydrolyzable ester thereof as well as metabolites of 9-cis retinoic 30

acid for the manufacture of a medicament for the treatment of T-helper cell type 1 (Th1) mediated immune diseases.

The invention also relates to a method for treating patients having T-helper cell type 1 (Th1) mediated immune diseases comprising administering to said human patient a compound selected from the group consisting of 9-cis retinoic acid, pharmaceutically acceptable salts and pharmaceutically acceptable hydrolyzable esters thereof, 9-cis retinal and pharmaceutically acceptable acetals thereof as well as 9-cis retinol and pharmaceutically acceptable hydrolyzable esters thereof as well as metabolites of 9-cis retinoic acid said compound being administered in an amount effective to treat said disease.

In the scope of the present invention, the term „T-helper cell type 1 mediated immune diseases“ relates to diseases with dominant cellular immune response, and it encompasses, in particular, autoimmune and other cell-mediated immune diseases, such as rheumatoid arthritis, multiple sclerosis, uveoretinitis, thyreoiditis, Crohn's disease, insulin dependent diabetes mellitus, eczema, systemic lupus erythematosus and allogeneic graft rejection (e.g. rejection of allogeneic skin, kidney, heart, liver or lung transplants). The term „eczema“ relates, in particular, to eczema due to delayed type hypersensitivity. The term „treatment“ or „treating“ includes preventive and/or therapeutic treatments.

9-cis retinoic acid and its derivatives and metabolic precursors and prodrugs as well as metabolites of 9-cis retinoic acid when administered to patients are effective, in particular in the therapy of the following T-helper cell

type 1 (Th1) mediated diseases: rheumatoid arthritis, multiple sclerosis, uveoretinitis, thyreoiditis, Crohn's disease, insulin dependent diabetes mellitus, systemic lupus erythematosus as well as eczema with its various
5 classes of exogenous eczema, such as irritant dermatitis, allergic contact dermatitis, tylotic eczema or pompholyx, endogenous eczema, such as seborrheic eczema (also called seborrhoic dermatitis or seborrheic dermatitis), asteatotic eczema and discoid eczema, and eczemas localised at various
10 sites of the body. 9-cis retinoic acid and its derivatives and metabolic precursors and prodrugs are effective in all those immune diseases which might be somehow linked with an increase of Th1 cell activity and an increased secretion of the related cytokines interleukin-12, interleukin-2,
15 interferon γ and tumor necrosis factor α , β .

For the treatment of Th1 mediated diseases other than eczema, the active compound, i.e. 9-cis retinoic acid, a pharmaceutically acceptable salt or a pharmaceutically acceptable hydrolyzable ester thereof, 9-cis retinal or a
20 pharmaceutically acceptable acetal thereof or 9-cis retinol or a pharmaceutically acceptable hydrolyzable ester thereof or metabolites of 9-cis retinoic acid, is administered orally. For the treatment of eczema the active compound is administered either orally or topically. Preferably, said
25 compound is administered as a composition containing said active compound and a pharmaceutically acceptable carrier or diluent compatible with said active compound. In preparing such composition, any conventional pharmaceutically acceptable carrier can be utilized. When
30 the drug is administered orally, it is generally administered at regular intervals, conveniently at mealtimes or once daily. It has been established that this

compound is effective in doses which show no or only mild side effects when given orally or when given topically. Therefore, oral administration of the active compound is generally preferred. For treating eczema however topical
5 administration may also be used advantageously.

In the treatment of T-helper cell type 1 mediated immune diseases, 9-cis retinoic acid and its derivatives and metabolic precursors and prodrugs as well as its metabolites, when administered orally, are therapeutically
10 efficacious in doses which induce no adverse events or only such mild side effect as dry lips and transient headache. At present, all retinoids exerting therapeutic effects in dermatological and oncological indications have to be administered orally in doses which induce more or less
15 marked side effects, belonging to the toxic syndrome of hypervitaminosis A, such as mucocutaneous, musculoskeletal and neurologic manifestations, particularly headache. In addition, they produce laboratory abnormalities such as elevated transaminases (ALAT, ASAT), elevated alkaline
20 phosphatase, as well as elevated triglycerides and cholesterol. In contrast, the daily doses of 9-cis retinoic acid and its derivatives and metabolic precursors and prodrugs and metabolites of 9-cis retinoic acid (typically 20 to 60 mg) therapeutically efficacious in T-helper cell
25 type 1 mediated immune diseases produce only very slight side effects, such as dry lips and transient headache, whereas all the other toxic signs and symptoms of the hypervitaminosis A syndrome, including the laboratory abnormalities, were not induced.

30 These same low daily doses of 9-cis retinoic acid, however, had no therapeutic effect on non-malignant skin-

disorders, such as acne, psoriasis, lamellar ichthyosis, Darier's disease and lichen planus. In summary, it was found that the very well tolerated low daily doses of 20 to 60 mg of 9-cis retinoic acid (and its derivatives and metabolic precursors and prodrugs) are efficacious in the treatment of T-helper cell type 1 mediated immune diseases, whereas such doses are not efficacious in the treatment of non-malignant skin disorders, such as acne, psoriasis and other keratinizing dermatoses. In malignant skin diseases and solid tumors of other organs, even high oral daily doses of 9-cis retinoic acid of up to 300 mg, inducing marked to severe side effects, did not lead to major objective tumor regressions.

In the treatment of T-helper cell type 1 mediated immune diseases, 9-cis retinoic acid, a pharmaceutically acceptable salt or a pharmaceutically acceptable hydrolyzable ester thereof, 9-cis retinal or a pharmaceutically acceptable acetal thereof or 9-cis retinol or a pharmaceutically acceptable hydrolyzable ester thereof or metabolites of 9-cis retinoic acid can be used alone or in combination with other measures, e.g. in combination with other pharmaceutically active substances such as topical or systemic corticosteroids and other immunosuppressive agents (cytostatics, antimetabolites, biological response modifiers, e.g. interferons, interleukins and other cytokines). If used in combination with other substances, 9-cis retinoic acid or its derivative or metabolic precursor or prodrug or its metabolites and said other substance can be administered separately or, preferably, incorporated in effective amounts into one pharmaceutical composition.

In the scope of the present invention, the „pharmaceutically acceptable salts“ includes any salt chemically permissible in the art for 9-cis retinoic acid and applicable to human patients in a pharmaceutically acceptable
5 preparation. Any such conventional pharmaceutically acceptable salt of 9-cis retinoic acid can be utilized. Among the conventional salts which can be utilized there are the base salts included, for example, alkali metal salts such as the sodium or potassium salt, alkaline earth metal salts such
10 as the calcium or magnesium salt, and ammonium or alkyl ammonium salts.

In accordance with this invention the 9-cis retinoic acid can also be administered in the form of its pharmaceutically acceptable hydrolyzable esters. Any pharmaceuti-
15 cally acceptable hydrolyzable ester can be used in the compositions and methods of this invention. Among the preferred esters are: the aromatic esters such as benzyl esters in which the benzyl moiety is unsubstituted or substituted with lower alkyl, halo, nitro, thio, or substituted
20 thio; or lower alkyl esters, e.g. ethyl, t-butyl, cyclopentyl, cyclohexyl or cycloheptyl ester; or 9-fluorenylmethyl ester.

In the scope of the present invention the term „alkyl“ means straight-chain, branched or cyclic alkyl residues, in particular those containing from 1 to 12 carbon
25 atoms, such as methyl, ethyl, propyl, isopropyl, t-butyl, decyl, dodecyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. The term „lower alkyl“ means alkyl groups containing from 1 to 7 carbon atoms.

In accordance with this invention a metabolic precursor or prodrug of 9-cis retinoic acid or metabolites of 9-cis retinoic acid are, in particular, 9-cis retinal, 9-cis retinol, a pharmaceutically acceptable acetal of 9-cis retinal or a pharmaceutically acceptable hydrolyzable esters of 9-cis retinol can alternatively be used instead of 9-cis retinoic acid, and any pharmaceutically acceptable acetal of 9-cis retinal and any pharmaceutically acceptable hydrolyzable ester of 9-cis retinol or 4-oxo-9-cis retinoic acid can be used in the compositions and methods of this invention. Among the preferred acetals of retinal are dialkyl acetals, especially di(lower alkyl) acetals such as the diethyl acetal, and dibenzyl acetals, wherein the benzyl moieties are unsubstituted or substituted with lower alkyl, halo, nitro, thio or substituted thio. Among the preferred hydrolyzable esters of 9-cis retinol are the esters formed with C₁-C₂₀-carboxylic acids such as C₁-C₂₀-alkanoic acids and C₁-C₂₀-alkenoic acids; particularly preferred are those carboxylic acid esters which contain and even number of carbon atoms in the carboxylic acid moiety such as acetate, stearate or palmitate.

The aforementioned 9-cis retinoic acid and its salts, its esters and its metabolic precursors or prodrugs as well as its metabolites are useful especially in pharmaceutically acceptable oral or topical modes. These pharmaceutical compositions contain said active compound in association with a compatible pharmaceutically acceptable carrier material. Any conventional carrier material can be utilized. The carrier material can be an organic or inorganic inert carrier material suitable for oral administration. Suitable carriers include water, gelatin, gum arabic, lactose, starch, magnesium stearate, talc,

vegetable oils, polyalkylene-glycols, petroleum jelly and the like. Furthermore, the pharmaceutically active preparations may contain other pharmaceutically active agents. Additionally, additives such as flavouring agents, 5 preservatives, stabilizers, emulsifying agents, buffers and the like may be added in accordance with accepted practices of pharmaceutical compounding.

The pharmaceutical preparations can be made up in any conventional form including inter alia: (a) a solid form 10 for oral administration such as tablets, capsules (e.g. hard or soft gelatine capsules), pills, sachets, powders, granules, and the like; and (b) preparations for topical administrations such as solutions, suspensions, ointments, creams, gels, micronized powders, aerosols and the like. 15 The pharmaceutical preparations may be sterilized and/or may contain adjuvants such as preservatives, stabilizers, wetting agents, emulsifiers, salts for varying the osmotic pressure and/or buffers.

For topical administration to the skin or mucous mem- 20 brane the aforementioned derivative is preferably prepared as ointments, tinctures, creams, gels, solution, lotions, sprays, suspensions, shampoos, hair soaps, perfumes and the like. In fact, any conventional composition can be utilized in this invention. Among the preferred methods of applying 25 the composition containing the agents of this invention is in the form of an ointment, cream or lotion. The pharmaceutical preparation for topical administration to the skin can be prepared by mixing the aforementioned active ingredient with non-toxic, therapeutically inert, solid or liquid 30 carriers customarily used in such preparation. These preparations generally contain at least about 0.0005 per-

cent by weight, preferably 0.0005 to 0.05 and more preferably about 0.001 to 0.01 percent by weight, of the active ingredient (i.e. 9-cis retinoic acid or its derivative or its metabolic precursor or prodrug or metabolites) based
5 upon the total weight of the composition. Since toxicity and irritancy of the active ingredient varies, depending on the kind of tissue - normal or pathologically altered - on which it is applied, it may however often be used in topical compositions in amounts up to 0.15 percent by
10 weight or even higher amounts. It is also preferred to apply these preparations once or twice daily to the skin. These preparations can be applied according to the need of the patient. In carrying out this invention, the active ingredient can be applied in an aqueous solution or an
15 alcohol solution such as ethanol.

In preparing the topical preparations described above, additives such as preservatives, thickeners, perfumes and the like conventional in the art of pharmaceutical compounding of topical preparation can be used. In addition,
20 conventional antioxidants or mixtures of conventional antioxidants can be incorporated into the topical preparations containing the aforementioned active agent. Among the conventional antioxidants which can be utilized in these preparations are included N-methyl- α -tocopherol-
25 amine, tocopherols, butylated hydroxyanisole, butylated hydroxytoluene, ethoxyquin and the like. Cream-base pharmaceutical formulations containing the active agent, used in accordance with this invention, are composed of aqueous emulsions containing a fatty acid alcohol, semi-solid petroleum hydrocarbon, ethylene glykol and an emulsifying
30 agent.

Ointment formulations containing the active agent in accordance with this invention comprise admixtures of a semi-solid petroleum hydrocarbon with a solvent dispersion of the active material. Cream compositions containing the
5 active ingredient for use in this invention preferably comprise emulsions formed from a water phase of a humectant, a viscosity stabilizer and water, an oil phase of a fatty acid alcohol, a semi-solid petroleum hydrocarbon and an emulsifying agent and a phase containing the active agent
10 dispersed in an aqueous stabilizer-buffer solution. Stabilizers may be added to the topical preparation. Any conventional stabilizer can be utilized in accordance with this invention. In the oil phase, fatty acid alcohol components function as a stabilizer. These fatty acid alcohol compo-
15 nents are derived from the reduction of a long-chain saturated fatty acid containing at least about 14 carbon atoms. Also, conventional perfumes and lotions generally utilized in topical preparation for the hair can be utilized in accordance with this invention. Furthermore, if desired, con-
20 ventional emulsifying agents can be utilized in the topical preparations of this invention.

A preferred oral dosage form comprises tablets, pills, sachets, or capsules of hard or soft gelatin, methylcellulose or of another suitable material easily dis-
25 solved in the digestive tract. Each tablet, pill, sachet or capsule can preferably contain from about 5 to about 50 mg, more preferably from about 10 to about 20 mg, of active ingredient. The oral dosages contemplated in accordance with the present invention will vary in accordance with the
30 needs of the individual patient as determined by the prescribing physician. Generally, however, a daily dosage of from about 0.05 mg to about 1.5 mg per kg of body weight

and preferably from about 0.3 mg to about 0.9 mg per kg of body weight of the patient is utilized. This dosage may be administered according to any dosage schedule determined by the physician in accordance with the requirements of the
5 patient.

Oral daily doses of from about 0.05 mg to about 1.5 mg per kg of body weight and preferably from about 0.3 mg to about 0.9 mg per kg of body weight are administered either continuously or on an intermittent
10 schedule e.g. in repetitive cycles of weekly 3 days on-, and 4 days off-treatment, or in cycles of alternatively 7 days on-, and 7 days off-treatment. For maintenance of a sufficiently high blood plasma or tissue level, it can be necessary to avoid concomitant medication of P450 isoenzyme
15 inducers and/or to add inhibitors of P450 isoenzymes. Dosage schedules are dependent on the kind of Th1-mediated immune disease such as e.g. eczema, rheumatoid arthritis, multiple sclerosis or Crohn's disease, but also on the stage of disease, plasma and tissue levels of 9-cis
20 retinoic acid, concomitant drug therapy and the patients' condition. 9-cis retinoic acid therapy can be given as continuous daily treatment or on an intermittent schedule as described above. 9-cis retinoic acid therapy can also consist in a continuous treatment as an induction therapy,
25 until a remission is achieved, and followed subsequently by a treatment on an intermittent schedule as a maintenance therapy.

9-cis retinoic acid treatment can be combined with other drugs used for treatment of Th1-mediated diseases,
30 such as e.g. rheumatoid arthritis, multiple sclerosis or Crohn's disease. Such drugs are e.g. methotrexate,

azathioprim, corticosteroids, cyclosporin, mycophenolic mofetil or interferons e.g. interferon β .

The dosage for treatment typically depends on the route of administration, the age, weight and disease condition of the individual. Suitable dosage forms are known in the art or can be easily obtained in a manner known per se. Formulations of lotions, gels, creams, hard or soft gelatin capsules, tablets and sachets that are particularly suitable in the scope of the present invention or that can be easily adjusted in accordance with the above teaching are disclosed e.g. in US-A-5,428,071.

In a preferred embodiment, the treatment of eczematoid conditions is carried out orally or topically.

In one embodiment, this invention provides a method of treating an eczematoid condition in a human patient afflicted with the condition, comprising orally administering or delivering to said patient 9-cis retinoic acid in an amount effective to treat the condition in said patient. A patient having active eczema is treated to lessen the severity of the eczema lesions or other symptoms. In the case where a patient previously had active eczema lesions, which have been brought into remission, this invention provides prophylactic treatment against the recurrence of active manifestations of eczema. In accordance with this invention, any eczematoid condition can be treated by the described method. Examples of the active or remission state eczematoid conditions which are treated include acute and chronic irritant dermatitis, allergic contact dermatitis, tylotic eczema or pompholyx, seborrhoic eczema, asteatotic eczema, and discoid eczema.

9-cis retinoic acid can be administered orally in any amount which is effective to treat the eczematoid condition. for example, the 9-cis retinoic acid or a precursor thereof which delivers from about 0.3 to about 5 0.9 milligrams of 9-cis retinoic acid per kilogram of body weight is preferred. This dosage may be administered according to any dosage schedule determined by the physician in accordance with the requirements of the patient. In accordance with specific embodiments, oral 10 daily doses from about 0.3 mg to about 0.9 mg per kg of body weight are administered either continuously or on an intermittent schedule e.g. in repetitive cycles of weekly 3 days on-, and 4 days off-treatment, or in cycles of alternatively 7 days on-, and 7 days off-treatment. 9-cis 15 retinoic acid therapy can also consist in a continuous treatment as an induction therapy, until a remission is achieved, and followed subsequently by a treatment on an intermittent schedule as a maintenance therapy.

For treatment of eczematoid conditions by oral delivery 20 of 9-cis retinoic acid, the 9-cis retinoic acid is delivered by administering orally to the patient a composition comprising 9-cis retinoic acid or a precursor thereof which delivers or produces 9-cis retinoic acid. Examples of suitable precursors of 9-cis retinoic acid 25 include pharmaceutically acceptable salts of 9-cis retinoic acid, hydrolyzable esters of 9-cis retinoic acid, 9-cis retinal, hydrolyzable esters 9-cis retinal, 9-cis retinol, and hydrolyzable esters of 9-cis retinol.

In an embodiment of the method of oral treatment of 30 eczematoid conditions, the composition containing 9-cis retinoic acid or precursor thereof is present in an oral

unit dosage form. In a more specific embodiment the oral unit dosage form is a tablet, capsule, pill or sachet containing from five to fifty milligrams, preferably from 10 to 20 mg of 9-cis retinoic acid or a pharmaceutically acceptable salt or hydrolyzable ester thereof.

The pharmaceutically acceptable salts include any salt chemically permissible for 9-cis retinoic acid and applicable to human patients in a pharmaceutically acceptable preparation. Any conventional pharmaceutically acceptable salt of 9-cis retinoic acid can be utilized. Among the conventional salts which can be utilized are included the base salts, for example, alkali metal salts such as sodium or potassium, alkaline earth metal salts such as calcium or magnesium, and ammonium or alkyl ammonium salts.

Pharmaceutically acceptable hydrolyzable esters can be used in the compositions and method of this invention. Among the esters are the aromatic esters such as benzyl (OBzl) or benzyl substituted with lower alkyl, halo, nitro, thio, or substituted thio, i.e., lower alkyl (1-7 carbon atoms) thio; aliphatic esters such as lower alkyl, t-butyl, cyclopentyl, cyclohexyl, cycloheptyl, and 9-fluorenylmethyl.

In another embodiment, this invention provides a method of treating an eczematoid condition of the skin in a human patient in need of such treatment, comprising topically applying to the area of the skin of said patient subject to the condition 9-cis retinoic acid in an amount effective to alleviate the effects of the condition. A patient having active eczema is treated to lessen the severity of the

eczema lesions or other symptoms. In the case where a patient previously had active eczema lesions, which have been brought into remission, this invention provides prophylactic treatment against the recurrence of active
5 manifestations of eczema. In accordance with this invention, any eczematoid condition can be treated by the described method. Examples of the active or remission state eczematoid conditions which are treated include irritant dermatitis, allergic contact dermatitis, tylotic eczema or
10 pompholyx, seborrheic eczema, asteatotic eczema, and discoid eczema.

For treatment of eczematoid conditions by topical delivery of 9-cis retinoic acid, the 9-cis retinoic acid is delivered by administering topically to the patient a
15 topical composition comprising a compound selected from 9-cis retinoic acid and dermatologically acceptable salts thereof. In formulating these topical compositions, the compound of formula I, its pharmaceutically acceptable salts or its pharmaceutically acceptable esters are
20 admixed with a pharmaceutically acceptable carrier for topical administration. Any conventional pharmaceutically acceptable carrier can be utilized in accordance with this invention.

These topical compositions which contain 9-cis
25 retinoic acid as well as its salts can contain any of the conventional excipients and additives commonly used in preparing topical compositions. Among the conventional additives or excipients which can be utilized in preparing these compositions in accordance with this invention are
30 preservatives, thickeners, perfumes and the like. In addition, the conventional antioxidants, such as butylated

hydroxyanisoles (BHA), ascorbyl palmitate, propyl gallate, citric acid, butylated hydroxytoluene (BHT), ethoxyquin, tocopherol, and the like can be incorporated into these compositions. These topical compositions can contain
5 conventional acceptable carriers for topical applications which are generally utilized in these compositions. These compositions may contain thickening agents, humectants, emulsifying agents and viscosity stabilizers, such as those generally utilized. In addition, these compositions
10 can contain colorants and perfumes which are conventional in preparing cosmetic compositions.

The composition for topical administration can contain any concentration of 9-cis retinoic acid or a salt thereof which is effective to treat the eczematoid condition. For
15 example, a composition which contains the compound in a concentration from about 0.001 to about 0.05 percent by weight of the composition can be used. More specific concentrations of the compound which are suitable in the method include from about 0.003 to about 0.03 percent by
20 weight of the composition; and from about 0.005 to about 0.01 percent by weight of the composition. Other suitable concentrations of the compound are at least about 0.0005 percent by weight; 0.0005 to 0.15 percent by weight; 0.0005 to 0.05 percent by weight; and 0.001 to 0.01
25 percent by weight of the composition.

The pharmaceutically acceptable salts include any salt chemically permissible for 9-cis retinoic acid and applicable to human patients in a pharmaceutically acceptable preparation. Any conventional pharmaceutically
30 acceptable salt of 9-cis retinoic acid can be utilized. Among the conventional salts which can be utilized are

included the base salts, for example, alkali metal salts such as sodium or potassium, alkaline earth metal salts such as calcium or magnesium, and ammonium or alkyl ammonium salts.

5 The topical composition of may be in any form which is conventional for topical compositions. For example, the topical composition can be present in the form of an ointment, cream, gel, lotion, or shampoo. Other examples of suitable forms for the topical composition include
10 tinctures, solutions, suspensions, micronized powders, suspensions, soaps, perfumes and aerosols.

The invention will be better understood from the following example, which is illustrative of the invention and not limiting.

15

Example 1

Activity of 9-cis retinoic acid in chronic hand eczema

a) Methods

Fifteen patients, eight men and seven women, with chronic hand eczema, refractory to conventional treatment,
20 were treated with 9-cis retinoic acid. Their mean age was 52.3 years, range 21-83. Before the start of 9-cis retinoic acid therapy, their eczema had already lasted for 3 months to 8 years, with a mean of 29 months. Besides avoidance of irritants and allergens, their previous treatment consisted
25 in topical topical corticosteroids (15 patients), topical tar (2 patients), isotretinoin (2 patients), tretinoin (1 patient) and X-rays (3 patients). The response to these treatments was in all cases unsatisfactory, in 4 patients moderate in 3 patients slight and in 8 patients no response

at all. Therapy consisted in a once daily oral dose of 40 mg 9-cis retinoic acid, given in the form of two soft gelatin capsules containing 20 mg of 9-cis retinoic acid each, with breakfast. 8 patients received in the first week only 20 mg. Mean duration of treatment was 2 months, range 1-3 months. The following lesions and symptoms were recorded on a 0-4 scale (0 = none, 1 = mild, 2 = moderate and 4 = severe) and used for evaluating the therapeutic effect: Erythema, papules and vesicles, desquamation, hyperkeratosis, rhagades and pruritus/pain. Side effects, particularly those belonging to the hypervitaminosis A syndrome, were recorded: Headache, dry lips, other mucocutaneous manifestations, muscoskeletal symptoms and laboratory abnormalities.

b) Results

As can be seen from Table 1, all fifteen patients responded markedly to 9-cis retinoic acid (9-cis-RA) and all the various lesions and symptoms were improved by the treatment. The total lesion-symptom score of the 14 responding patients was reduced by a mean of 81.3% (range 53-100 %). The various lesions and symptoms such as erythema, papules and vesicles etc., were all favourably influenced and regressed by 62-100 %. 9-cis retinoic acid in a dose of 40 mg daily was very well tolerated. The only side effect noted in these 15 patients were transient headache in 2, and dry lips in 5 patients. No other mucocutaneous manifestations, seen with higher doses and no musculoskeletal or other symptoms were observed. Such symptoms were seen with higher doses [Kurie et al., Clin. Cancer Res. 2, 287-293 (1996); Miller et al., Clin. Cancer Res. 2, 471-475 (1996)]. The well known laboratory abnormalities, such as elevation of transaminases (ALAT, ASAT),

alkaline phosphatase, triglycerides and cholesterol, frequently caused by retinoids were not seen with this low dosage of 9-cis retinoic acid. The response to the therapy with 9-cis retinoic acid in patients chronic hand eczema, s refractory to conventional treatment, was assessed by the doctor, as well as by the patient and was considered as very good or good in 13 of 15 patients or 87 % of the patients (Tables 1 and 2).

Table 1 : Treatment of Chronic Hand Eczema with 9-cis Retinoic Acid

Patient No.	1	2	3	4	5
Sex	f	m	m	m	m
Age (years)	83	41	49	50	37
Duration of eczema (months)	3	20	16	60	36
Previous treatment	top. steroids	top. steroids	top. steroids x-rays	top. steroids	top. steroids x-rays
Response to previous treatment	no	slight	moderate	moderate	moderate
9-cis RA					
Oral daily dose (mg)	40	40	40	40	40
Duration of treatment (months)	2	2	2	3	1
Efficacy (0-4 scale) before / after treatment					
Erythema	1 / 0	1 / 0	1 / 0.5	1 / 0.5	2 / 0
Papules, vesicles	3 / 0	3 / 1	2 / 1	2 / 1	2 / 0
Pruritus (+ pain)	2 / 0	2 / 0	3 / 1	3 / 1	3 / 2
Desquamation	2 / 1	3 / 1	3 / 1	3 / 1	2 / 1
Hyperkeratosis	4 / 0	3 / 0	1 / 1	2 / 1	1 / 0
Rhagades	3 / 0	2 / 0	1 / 0	0.5 / 0	1 / 0
Total Score (0-4 scale)	15 / 1	14 / 2	11 / 4.5	11 / 4.5	11 / 3
Reduction of score in %	93	86	59	61	73
Response assessed by patient	very good	very good	good	very good	very good
Response assessed by doctor	very good	very good	good	good	good

Table 1 : Treatment of Chronic Hand Eczema with 9-cis Retinoic Acid (continued)

Patient No.	6	7	8	9	10
Sex	m	f	m	m	f
Age (years)	43	78	62	53	51
Duration of eczema (months)	24	36	96	6	48
Previous treatment	top. steroids isotretinoin tretinoin, acitretin	top. steroids x-rays	top. steroids	top. steroids	top. steroids isotretinoin
Response to previous treatment	no	no	slight	no	no
9-cis RA					
Oral daily dose (mg)	40	40	40	40	40
Duration of treatment (months)	2	3	2	2.5	1.5
Efficacy (0-4 scale) before / after treatment					
Erythema	1 / 0	2 / 0	4 / 2	0 / 0	2 / 2
Papules, vesicles	1 / 0	0 / 0	3 / 1	0 / 0	0 / 0
Pruritus (+ pain)	4 / 0	5 / 0	5 / 2	4 / 1	7 / 7
Desquamation	1 / 0	3 / 0	1 / 0	1 / 0	3 / 3
Hyperkeratosis	2 / 0	1 / 0	1 / 1	2 / 0	3 / 3
Rhagades	1 / 0	3 / 0	3 / 2	2 / 0	2 / 2
Total Score (0-4 scale)	10 / 0	14 / 0	17 / 8	9 / 1	17 / 17
Reduction of score in %	100	100	53	89	0
Response assessed by patient	very good	very good	moderate	very good	no
Response assessed by doctor	very good	very good	good	very good	no

Table 1 : Treatment of Chronic Hand Eczema with 9-cis Retinoic Acid (continued)

Patient No.	11	12	13	14	15
Sex	f	f	f	m	f
Age (years)	66	49	50	51	21
Duration of eczema (months)	48	12	48	18	8
Previous treatment	top. steroids	top. steroids	top. steroids tar	top. steroids	top. steroids
Response to previous treatment	slight	no	no	no	moderate
9-cis RA					
Oral daily dose (mg)	40	40	40	40	40
Duration of treatment (months)	1.5	2	2.5	1	2
Efficacy (0-4 scale) before / after treatment					
Erythema	2/0	1/0	1/0	2/1	1/0
Papules, vesicles	1/0	0/0	0/0	0/0	1/0.5
Pruritus (+ pain)	4/0	5/0	5/0.5	3/1	2/0.5
Desquamation	2/0	0/0	0/0	2/0	1/0.5
Hyperkeratosis	2/1	2/0	2/0.5	2/0	2/0.5
Rhagades	2/1	2/0	1/0	2/0	1/0
Total Score (0-4 scale)	13/2	10/0	9/1	11/2	8/2
Reduction of score in %	85	100	89	82	75
Response assessed by patient	very good	very good	good	very good	good
Response assessed by doctor	good	very good	good	good	moderate

Table 2 : Therapy with oral 9-cis Retinoic Acid (9-cis RA) in Chronic Hand Eczema (Summary)

No. of patients	15
Sex	8 males, 7 females
Age	21 - 83 years, mean 52.3 years
Duration of eczema	3 - 96 months, mean 29 months
Previous treatment	topical corticosteroids (15), tar (2), isotretinoin (2), tretinoin (1), acitretin (1)
Response to previous treatment	moderate (4), slight (3), no (8)
9-cis RA Therapy	
Daily oral dose	40 mg
Duration of treatment	1 - 3 months, mean 2 months
Efficacy	
Reduction of Lesion / symptom score, in %	for all 15 patients 0-100, mean 76.3 for the 14 responding patients 53 -100, mean 81.8

Response of patients to 9-cis RA therapy

Assessment by	Patient	Doctor
very good	10 (67%)	6 (40%)
good	3 (20%)	7 (47%)
moderate	1 (6.5%)	1 (6.5%)
slight	0 (0%)	0 (0%)
no	1 (6.5%)	1 (6.5%)
Side effects		
Dry lips	5 (33.3%)	
Headache, transient	2 (13.3%)	

Claims

1. The use of 9-cis retinoic acid and pharmaceutically acceptable salts and pharmaceutically acceptable hydrolyzable esters thereof as well as prodrugs and
5 metabolites thereof and their pharmaceutically acceptable salts and pharmaceutically acceptable hydrolyzable esters for the manufacture of a medicament for the treatment of T-helper cell type 1 mediated immune diseases.
2. The use of a compound selected from the group consisting of 9-cis retinoic acid and pharmaceutically acceptable
10 salts and pharmaceutically acceptable hydrolyzable esters thereof, 9-cis retinal and pharmaceutically acceptable acetals thereof, and 9-cis retinol and pharmaceutically acceptable hydrolyzable esters thereof as active ingredient
15 for the manufacture of a medicament for the treatment of T-helper cell type 1 mediated immune diseases.
3. The use according to claim 1 or 2, wherein the active ingredient is used in combination with a pharmaceutically acceptable carrier.
- 20 4. The use according to claim 1, 2 or 3, wherein the medicament is manufactured for oral or topical administration.
5. The use according to any one of claims 1 to 4, wherein the medicament is manufactured as a tablet, capsule, pill,
25 sachet, ointment, cream or lotion.

6. The use according to any one of claims 1 to 5, wherein the medicament is manufactured as a tablet, capsule, pill or sachet containing 5 to 50 mg, preferably 10 to 20 mg, of active ingredient.
- 5 7. The use according to any one of claims 1 to 6, wherein the medicament is manufactured for oral daily dosage of from 0.05 mg to 1.5 mg, preferably from 0.3 mg to 0.9 mg, per kg of body weight.
8. The use according to any one of claims 1 to 5, wherein
10 the medicament is manufactured as an ointment, cream or lotion containing 0.0005 to 0.05 percent by weight, preferably 0.001 to 0.01 percent by weight, of the active ingredient.
9. The use according to any one of claims 1 to 8, wherein
15 the active ingredient is selected from the group consisting of 9-cis retinoic acid and alkali metal salts, alkaline earth metal salts, benzyl esters, lower alkyl esters and 9-fluorenylmethyl esters thereof, 9-cis retinal and dialkyl acetals and dibenzyl acetal thereof, and 9-cis retinol and
20 esters thereof formed with C₁-C₂₀-carboxylic acids.
10. The use according to claims 1 to 9 wherein the active ingredient is 9-cis retinoic acid or a pharmaceutically acceptable salt thereof.
11. The use according to any one of claims 1 to 7, 9 and
25 10, wherein the medicament is manufactured for the treatment of autoimmune diseases, and other cell-mediated immune diseases such as rheumatoid arthritis, multiple sclerosis, uveoretinitis, thyroiditis, Crohn's disease,

insulin dependent diabetes mellitus, eczema, systemic lupus erythematosus or allogeneic graft rejection.

12. The use according to any one of claims 1 to 10, wherein the medicament is manufactured for the treatment of exogenous eczema, such as irritant dermatitis, allergic contact dermatitis, tylotic eczema, pompholyx or of endogenous eczema, such as seborrheic eczema, asteatotic eczema and discoid eczema.

13. A method of treating an eczematoid condition in a human patient afflicted with the condition, comprising orally administering to said patient 9-cis retinoic acid in an amount effective to treat the condition in said patient.

14. The method of claim 13, wherein the condition comprises active eczema lesions.

15. The method of claim 14, wherein the condition comprises active eczema lesions of: irritant dermatitis, allergic contact dermatitis, seborrheic eczema, asteatotic eczema, or discoid eczema.

16. The method of claim 13, wherein the condition comprises preexisting eczema lesions which are in remission.

17. The method of claim 16, wherein the condition comprises preexisting lesions of: irritant dermatitis, allergic contact dermatitis, tylotic eczema or pompholyx, seborrheic dermatitis, asteatotic eczema, or discoid eczema; which lesions are in remission.

18. The method of claim 13, wherein the effective amount of the 9-cis retinoic acid is from about 0.3 to about 0.9 milligrams per kilogram of body weight.

19. The method of claim 13, wherein the 9-cis retinoic acid
5 is administered in a composition comprising 9-cis retinoic acid or a precursor thereof which delivers 9-cis retinoic acid.

20. The method of claim 19, wherein the precursor is selected from the group consisting of pharmaceutically
10 acceptable: salts of 9-cis retinoic acid; hydrolyzable esters of 9-cis retinoic acid; 9-cis retinal; hydrolyzable esters 9-cis retinal; 9-cis retinol; and hydrolyzable esters of 9-cis retinol.

21. The method of claim 19, wherein the composition is
15 present in an oral unit dosage form.

22. The method of claim 21, wherein the oral unit dosage form is a tablet, capsule, pill or sachet containing from twenty to sixty milligrams of 9-cis retinoic acid or a pharmaceutically acceptable salt or hydrolyzable ester
20 thereof.

23. A method of treating an eczematoid condition of the skin in a human patient in need of such treatment, comprising topically applying to the area of the skin of said patient subject to the condition 9-cis retinoic acid
25 in an amount effective to alleviate the effects of the condition.

24. The method of claim 23, wherein the condition comprises active eczema lesions.

25. The method of claim 24, wherein the condition comprises active eczema lesions of: irritant dermatitis, allergic
s contact dermatitis, tylotic eczema or pompholyx, seborrheic eczema, asteatotic eczema, or discoid eczema.

26. The method of claim 23, wherein the condition comprises preexisting eczema lesions which are in remission.

27. The method of claim 26, wherein the condition comprises
10 preexisting lesions of: irritant dermatitis, allergic contact dermatitis, tylotic eczema or pompholyx, seborrheic dermatitis, asteatotic eczema, or discoid eczema; which lesions are in remission.

28. The method of claim 23, wherein the 9-cis retinoic acid
15 is applied in a topical composition comprising a compound selected from the group consisting of 9-cis retinoic acid and dermatologically acceptable salts thereof.

29. The method of claim 28, wherein the composition contains the compound in an amount from about 0.001 to
20 about 0.05 percent by weight of the composition.

30. The method of claim 29, wherein the composition contains the compound in an amount from about 0.003 to about 0.03 percent by weight of the composition.

31. The method of claim 30, wherein the composition
25 contains the compound in an amount from about 0.005 to about 0.01 percent by weight of the composition.

- 30 -

32. The method of claim 31, wherein the composition is present in the form of an ointment, cream, gel, lotion, or shampoo.

33. A method of treating a T-helper cell type 1 mediated
5 immune disease in a human patient, comprising administering to the patient a compound in an amount effective to treat the disease; wherein the compound is selected from the group consisting of 9-cis retinoic acid, pharmaceutically acceptable salts and hydrolyzable esters thereof,
10 precursors and metabolites of 9-cis retinoic acid and pharmaceutically acceptable salts and hydrolyzable esters thereof.

34. The method of claim 33, wherein 9-cis retinoic acid or a pharmaceutically acceptable salt thereof is administered.

15 35. The method of claim 33, wherein a precursor of 9-cis retinoic acid selected from the group consisting of 9-cis retinal, pharmaceutically acceptable acetals of 9-cis retinal, 9-cis retinol, pharmaceutically acceptable hydrolyzable esters of 9-cis retinol, and pharmaceutically
20 acceptable salts thereof, is administered.

36. The method of claim 33, wherein the compound is administered orally.

37. The method of claim 36, wherein the effective amount of the compound is from 0.05 to 1.5 milligrams per kilograms
25 of body weight per day.

38. The method of claim 37, wherein effective amount of the compound is from 0.3 to 0.9 milligrams per kilograms of body weight per day.

39. The method of claim 33, wherein the disease is an
5 eczematoid condition and the compound is administered topically.

40. The method of claim 33, wherein the compound is administered in a composition comprising the compound and a pharmaceutically acceptable carrier.

10 41. The method of claim 40, wherein the composition is in the form of a tablet, capsule, pill or sachet.

42. The method of claim 41, each tablet, capsule, pill or sachet contains from five to fifty milligrams of the compound.

15 43. The method of claim 42, wherein each tablet, capsule, pill or sachet contains from ten to twenty milligrams of the compound.

44. The method of claim 40, wherein the composition is in the form of an ointment, cream or lotion.

20 45. The method of claim 44, wherein the composition contains the compound in an amount from 0.0005 to 0.05 percent by weight of the composition.

46. The method of claim 45, wherein the composition contains the compound in an amount from 0.001 to 0.01
25 percent by weight of the composition.

47. The method of claim 33, wherein the compound is selected from the group consisting of 9-cis retinoic acid and alkali metal salts, alkaline earth metal salts, benzyl esters, lower alkyl esters and 9-fluorenylmethyl esters thereof,
5 9-cis retinal and dialkyl acetals and dibenzyl acetal thereof, and 9-cis retinol and esters thereof formed with C₁-C₂₀-carboxylic acids.

48. The method of claim 36, wherein the disease which is treated is rheumatoid arthritis, multiple sclerosis,
10 uveoretinitis, thyreoiditis, Crohn's disease, insulin dependent diabetes mellitus, eczema, systemic lupus erythematosus or allogeneic graft rejection.

49. The method of claim 33, 36 or 39, wherein the disease is an exogenous eczema.

15 50. The method of claim 49, wherein the exogenous eczema is irritant dermatitis, allergic contact dermatitis, tylotic eczema or pompholyx.

51. The method of claim 33, 36 or 39, wherein the disease is an endogenous eczema.

20 52. The method of claim 51, wherein the endogenous eczema is seborrheic eczema, asteatotic eczema or discoid eczema.

* * * * *

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/05236

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/07 A61K31/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 22818 A (PFIZER) 13 October 1994 see page 4, line 13-14 ---	1,2,33
X	EP 0 579 915 A (HOFFMANN LA ROCHE) 26 January 1994 see page 2, line 1-12 see page 3, line 56 - page 4, line 5 ---	1-52
X	US 5 093 360 A (YU RUEY J ET AL) 3 March 1992 see column 1, line 41 - column 2, line 32 see column 3, line 56-58 ---	1-52
X	US 5 658 949 A (AGGARWAL BHARAT B) 19 August 1997 see column 5, line 35-40; claims 2,6,15 see column 5, line 58-61 ---	1-52
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

4 December 1998

Date of mailing of the international search report

11/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel.: (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Engl, 8

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/05236

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>YILI YANG ET AL.: "9-cis-Retinoic Acid inhibits activation-driven T-cell apoptosis: Implications for Retinoid X Receptor involvement in thymocyte development"</p> <p>PROC. NATL. ACAD. SCI. USA, vol. 90, July 1993, pages 6170-6174, XP000578397</p> <p>see abstract</p> <p>-----</p>	1-52

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern 1al Application No

PCT/EP 98/05236

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9422818	A	13-10-1994	NONE	
EP 0579915	A	26-01-1994	AU 3716193 A	25-11-1993
			CA 2096196 A	21-11-1993
			CN 1080525 A	12-01-1994
			HU 66864 A	30-01-1995
			JP 6032740 A	08-02-1994
			NO 931831 A	22-11-1993
			NZ 247630 A	21-12-1995
			AT 141914 T	15-09-1996
			AU 3863593 A	25-11-1993
			CN 1085211 A	13-04-1994
			DE 69304279 D	02-10-1996
			DK 580968 T	14-10-1996
			EP 0580968 A	02-02-1994
			ES 2091515 T	01-11-1996
			GR 3021833 T	28-02-1997
			JP 2670005 B	29-10-1997
			JP 6040975 A	15-02-1994
			MX 9302886 A	01-11-1993
			ZA 9303356 A	22-11-1993
			ZA 9303357 A	22-11-1993
US 5093360	A	03-03-1992	AU 649605 B	02-06-1994
			AU 4914690 A	11-10-1990
			CA 2009835 A	07-10-1990
			DE 69027057 D	27-06-1996
			DE 69027057 T	16-01-1997
			EP 0391033 A	10-10-1990
			US 5492935 A	20-02-1996
US 5658949	A	19-08-1997	US 5457129 A	10-10-1995
			AU 697072 B	24-09-1998
			AU 6944194 A	12-12-1994
			CA 2162990 A	24-11-1994
			EP 0708648 A	01-05-1996
			JP 8510727 T	12-11-1996
			WO 9426277 A	24-11-1994
			ZA 9403320 A	16-11-1995